

Pay as little as \$10 for TRINTELLIX with the Savings Card

If you are eligible,* you could
pay as little as \$10 for either a
30-day or 90-day prescription

Pay as little as \$10*

*Download card to see eligibility and term & conditions.

Save up to **\$100** on a 30-day or **\$300** on a 90-day prescription.

ACTIVATE your card at Trintellix.com/tAccess

RxBIN: 610524

RxPCN: Loyalty

RxGRP: **50776825**

ISSUER: (80840)

ID: **XXXXXXXXXX**

Please see accompanying Full Prescribing Information and Medication Guide, including Boxed WARNING for Suicidal Thoughts and Actions, and discuss with your doctor.

Powered By
McKesson

Trintellix
vortioxetine
5mg•10mg•20mg tablets



The TRINTELLIX Savings Card is available for adults with commercial health insurance taking TRINTELLIX. Savings of up to \$100 per 30-day or \$300 per 90-day prescription, with total savings of up to \$1,300 over one year. Restrictions apply.*

Check your eligibility and enroll today.

GET STARTED

*See the TRINTELLIX Savings Card for full Eligibility Requirements and Terms & Conditions. The Savings Card cannot be used by patients in federal-, state-, or government-funded healthcare programs, or by cash patients.

What is TRINTELLIX (vortioxetine)?

TRINTELLIX is a prescription medicine used in adults to treat a certain type of depression called Major Depressive Disorder (MDD). TRINTELLIX has not been shown to be safe and effective for use in children.

IMPORTANT SAFETY INFORMATION

Suicidal Thoughts & Actions

- TRINTELLIX and other antidepressants increase the risk of suicidal thoughts and actions in people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed.
- TRINTELLIX is not for use in children.
- Call your doctor or get emergency help right away if you have new or sudden changes in mood, behavior, thoughts or feelings, if you develop suicidal thoughts or actions, or if you have or develop symptoms that are new, worse, or worry you.

Please see additional Important Safety Information throughout, including Full Boxed WARNING for Suicidal Thoughts and Actions on page 4 and accompanying Medication Guide on pages 7 and 8, and discuss with your doctor.

Trintellix
vortioxetine
5mg•10mg•20mg tablets



TRINTELLIX may cost less than you think

TRINTELLIX is covered for 85% of patients in the US with commercial insurance

Cost is an important consideration after finding the right treatment. If you have commercial health insurance that covers medications, you may be eligible for the TRINTELLIX Savings Card, which could help you keep prescription costs down.*

QUICKLY CHECK YOUR ELIGIBILITY NOW

Let's get your Savings Card. It's easy!

If you are eligible for the Savings Card, you can get it and start using it today.



1. Check your eligibility

- Visit the [TRINTELLIX website](#) to see if you're eligible



2. Provide your contact information

- On the website, enter your contact information
- Download and print your Savings Card
 - You can also save it to your mobile wallet on your phone



3. Use your Savings Card

- Show your Savings Card to your pharmacist to start saving on your TRINTELLIX prescription

*Restrictions apply. See the Savings Card for Eligibility Requirements and Terms & Conditions. Formulary data provided by Fingertip Formulary® and current as of September 2022. Check your individual health plan for coverage and cost information.

IMPORTANT SAFETY INFORMATION

Who should not take TRINTELLIX?

Do not start or take TRINTELLIX if you:

- are allergic to vortioxetine or any of the ingredients in TRINTELLIX
- are taking, or have stopped taking within the last 14 days, a medicine called a Monoamine Oxidase Inhibitor (MAOI), including the antibiotic linezolid or intravenous methylene blue

Do not start taking an MAOI for at least 21 days after you stop treatment with TRINTELLIX.

Please see additional Important Safety Information throughout, including Full Boxed WARNING for Suicidal Thoughts and Actions on page 4 and accompanying Medication Guide on pages 7 and 8, and discuss with your doctor.

You're not alone. We've got you covered.

The tAccess Support Program is available for everyone

tAccess is a free support program designed to partner with you on your TRINTELLIX journey. It costs nothing to join and offers customized resources and tools. You can utilize this program, whether or not you qualify for the Savings Card.

Support tailored to your needs



INFORMATIVE EMAILS

Receive helpful tools, such as discussion guides for conversations with your doctor, as well as inspiring stories about MDD from real TRINTELLIX patients



INSPIRING TEXTS AND REMINDERS FOR DOSING AND REFILLS

Sign up to receive customized reminders and inspiring texts to help you stay on track with treatment. It can help you easily set up dosing and refill reminders on your phone



FREE CHATS WITH YOUR tACCESS ADVISOR

Have a question about support tools? Need help activating or replacing your Savings Card? Our support advisors are here to assist you and provide helpful resources and tips:

1-844-654-TRIN, (8AM–8PM EST, M–F)

tAccess is here to help. Visit [TRINTELLIX.com/tAccess](https://www.trintellix.com/tAccess) to learn more.

IMPORTANT SAFETY INFORMATION (cont'd)

What should I tell my doctor before taking TRINTELLIX?

Before taking TRINTELLIX, tell your doctor:

- about all your medical and other health conditions
- if you are pregnant or plan to become pregnant, since TRINTELLIX may harm your unborn baby. Taking TRINTELLIX during your third trimester may cause your baby to have withdrawal symptoms after birth or to be at increased risk for a serious lung problem at birth. Tell your doctor right away if you become or think you are pregnant while taking TRINTELLIX
- if you are breastfeeding or plan to breastfeed, since it is not known if TRINTELLIX passes into your breast milk

Please see additional Important Safety Information throughout, including Full Boxed WARNING for Suicidal Thoughts and Actions on page 4 and accompanying Medication Guide on pages 7 and 8, and discuss with your doctor.

What is TRINTELLIX (vortioxetine)?

TRINTELLIX is a prescription medicine used in adults to treat a certain type of depression called Major Depressive Disorder (MDD). TRINTELLIX has not been shown to be safe and effective for use in children.

IMPORTANT SAFETY INFORMATION

Suicidal Thoughts & Actions

TRINTELLIX and other antidepressants increase the risk of suicidal thoughts and actions in people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed. TRINTELLIX is not for use in children. Depression or other mental illnesses are the most important causes of suicidal thoughts or actions. Pay close attention to any changes, especially new or sudden changes in mood, behavior, thoughts, or feelings, or if you develop suicidal thoughts or actions. Call your doctor or get emergency help right away to report any of these, or if you have symptoms such as suicidal thoughts or actions, impulsivity, aggressive or violent actions, depression, anxiety or panic attacks, agitation, restlessness, anger, irritability, trouble sleeping, an increase in activity or talking, or other unusual changes in behavior or mood; especially if they are new, worse, or worry you.

Who should not take TRINTELLIX?

Do not start or take TRINTELLIX if you:

- are allergic to vortioxetine or any of the ingredients in TRINTELLIX
- are taking, or have stopped taking within the last 14 days, a medicine called a Monoamine Oxidase Inhibitor (MAOI), including the antibiotic linezolid or intravenous methylene blue

Do not start taking an MAOI for at least 21 days after you stop treatment with TRINTELLIX.

What should I tell my doctor before taking TRINTELLIX?

Before taking TRINTELLIX, tell your doctor:

- about all your medical and other health conditions
- if you are pregnant or plan to become pregnant, since TRINTELLIX may harm your unborn baby. Taking TRINTELLIX during your third trimester may cause your baby to have withdrawal symptoms after birth or to be at increased risk for a serious lung problem at birth. Tell your doctor right away if you become or think you are pregnant while taking TRINTELLIX
- if you are breastfeeding or plan to breastfeed, since it is not known if TRINTELLIX passes into your breast milk

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements, since TRINTELLIX and some medicines may cause serious side effects (or may not work as well) when taken together. **Especially tell your doctor if you take:** medicines for migraine headache called triptans; tricyclic antidepressants; lithium; tramadol, fentanyl, meperidine, methadone, or other opioids; tryptophan; buspirone; St. John's Wort; medicines that can affect blood clotting such as aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), warfarin; diuretics; medicines used to treat mood, anxiety, psychotic, or thought disorders, including selective serotonin reuptake inhibitors (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs); or medicines used to treat seizures or convulsions.

Please see additional Important Safety Information throughout, including Full Boxed WARNING for Suicidal Thoughts and Actions on page 4 and accompanying Medication Guide on pages 7 and 8, and discuss with your doctor.

What are the possible side effects of TRINTELLIX?

TRINTELLIX may cause serious side effects, including:

- **Serotonin syndrome:** A potentially life-threatening problem that can happen when you take TRINTELLIX with certain other medicines. Call your doctor or go to the nearest emergency room right away if you have any of the following signs and symptoms of serotonin syndrome: agitation; seeing or hearing things that are not real; confusion; coma; fast heart-beat; changes in blood pressure; dizziness; sweating; flushing; high body temperature; shaking, stiff muscles, or muscle twitching; loss of coordination; seizures; nausea, vomiting, diarrhea.
- **Increased risk of bleeding:** Taking TRINTELLIX with aspirin, NSAIDs, warfarin or blood thinners may add to this risk. Tell your doctor right away about any unusual bleeding or bruising.
- **Manic episodes:** Manic episodes may happen in people with bipolar disorder who take TRINTELLIX. Symptoms may include: greatly increased energy; racing thoughts; unusually grand ideas; talking more or faster than usual; severe problems sleeping; reckless behavior; excessive happiness or irritability.
- **Discontinuation syndrome:** Suddenly stopping TRINTELLIX may cause you to have serious side effects including: nausea; sweating; changes in your mood; irritability and agitation; dizziness; electric shock feeling; tremor; anxiety; confusion; headache; tiredness; problems sleeping; hypomania; ringing in your ears; seizures.
- **Eye problems:** TRINTELLIX may cause a type of eye problem called angle-closure glaucoma in people with certain other eye conditions. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are. Call your doctor if you have eye pain, changes in your vision, or swelling or redness in or around the eye.
- **Low levels of salt (sodium) in your blood:** Low sodium levels in your blood that may be serious and may cause death can happen during treatment with TRINTELLIX. Elderly people and people who take certain medicines may be at a greater risk for developing low sodium levels in their blood. Signs and symptoms may include headache; difficulty concentrating; memory changes; confusion; weakness and unsteadiness on your feet which can lead to falls. **In more severe or more sudden cases, signs and symptoms include:** seeing or hearing things that are not real; fainting; seizures; coma; stopping breathing.
- **Sexual problems:** Taking TRINTELLIX may cause sexual problems. Symptoms in males may include: delayed ejaculation or inability to have an ejaculation, decreased sex drive, or problems getting or keeping an erection. Symptoms in females may include: decreased sex drive, or delayed orgasm or inability to have an orgasm. Talk to your doctor if you develop any changes in your sexual function or if you have any questions or concerns about sexual problems during treatment with TRINTELLIX.

The most common side effects of TRINTELLIX include:

- nausea
- constipation
- vomiting

These are not all the possible side effects of TRINTELLIX. Tell your doctor if you have any side effect that bothers you or does not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout, including Full Boxed WARNING for Suicidal Thoughts and Actions on page 4 and accompanying Medication Guide on pages 7 and 8, and discuss with your doctor.

YOU MAY BE ELIGIBLE TO SAVE ON YOUR PRESCRIPTION

If you are eligible, you could pay as little as \$10 for either a 30-day or 90-day prescription with the TRINTELLIX Savings Card*

Text "INFO" to 26785 to learn more

*Restrictions apply. See the TRINTELLIX Savings Card for Eligibility Requirements and Terms & Conditions. Commercially insured patients only.

IMPORTANT SAFETY INFORMATION

Suicidal Thoughts & Actions

- TRINTELLIX and other antidepressants increase the risk of suicidal thoughts and actions in people 24 years of age and younger, especially within the first few months of treatment or when the dose is changed.
- TRINTELLIX is not for use in children.
- Call your doctor or get emergency help right away if you have new or sudden changes in mood, behavior, thoughts or feelings, if you develop suicidal thoughts or actions, or if you have or develop symptoms that are new, worse, or worry you.

Please see additional Important Safety Information throughout and accompanying Medication Guide on pages 7 and 8, and discuss with your doctor.


Trintellix
vortioxetine
5mg•10mg•20mg tablets



Visit Trintellix.com

TRINTELLIX is a trademark of H. Lundbeck A/S registered with the U.S. Patent and Trademark Office and used under license by Takeda Pharmaceuticals America, Inc.
TAKEDA and the TAKEDA logo are registered trademarks of Takeda Pharmaceutical Company Limited.
©2023 Takeda Pharmaceuticals U.S.A., Inc. All rights reserved. 1-877-TAKEDA-7 (1-877-825-3327).
US-VOR-1168v2.0 09/23

MEDICATION GUIDE

TRINTELLIX (trin'-TELL-ix) (vortioxetine) Tablets

What is the most important information I should know about TRINTELLIX?

TRINTELLIX can cause serious side effects, including:

- **Increased risk of suicidal thoughts and actions.** TRINTELLIX and other antidepressant medicines increase the risk of suicidal thoughts and actions in people 24 years of age and younger, **especially within the first few months of treatment or when the dose is changed. TRINTELLIX is not for use in children.**
 - Depression or other mental illnesses are the most important causes of suicidal thoughts or actions.

How can I watch for and try to prevent suicidal thoughts and actions?

- Pay close attention to any changes, especially sudden changes in mood, behavior, thoughts, or feelings, or if you develop suicidal thoughts or actions. This is very important when an antidepressant medicine is started or when the dose is changed.
- Call your healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings or if you develop suicidal thoughts or actions.
- Keep all follow-up visits with your healthcare provider as scheduled. Call your healthcare provider between visits as needed, especially if you have concerns about symptoms.

Call your healthcare provider or get emergency help right away if you have any of the following symptoms, especially if they are new, worse, or worry you:

- attempts to commit suicide
- thoughts about suicide or dying
- feeling agitated, restless, angry or irritable
- other unusual changes in behavior or mood
- acting on dangerous impulses
- new or worse depression
- trouble sleeping
- acting aggressive or violent
- new or worse anxiety or panic attacks
- an increase in activity or talking more than what is normal for you

What is TRINTELLIX?

TRINTELLIX is a prescription medicine used in adults to treat a certain type of depression called Major Depressive Disorder (MDD). TRINTELLIX has not been shown to be safe and effective for use in children.

Do not take TRINTELLIX if you:

- are allergic to vortioxetine or any of the ingredients in TRINTELLIX. See the end of this Medication Guide for a complete list of ingredients in TRINTELLIX
- are taking, or have stopped taking within the last 14 days, a medicine called a Monoamine Oxidase Inhibitor (MAOI), including the antibiotic linezolid or intravenous methylene blue

Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid or intravenous methylene blue.

Do not start taking an MAOI for at least 21 days after you stop treatment with TRINTELLIX.

Before taking TRINTELLIX, tell your healthcare provider about all of your medical conditions, including if you:

- have or had bleeding problems
- have, or have a family history of, bipolar disorder, mania, or hypomania
- have high pressure in the eye (glaucoma)
- have low sodium levels in your blood

- are pregnant or plan to become pregnant. TRINTELLIX may harm your unborn baby. Taking TRINTELLIX during your third trimester of pregnancy may cause your baby to have withdrawal symptoms after birth or may cause your baby to be at increased risk for a serious lung problem at birth. Talk to your healthcare provider about the risks to you and your unborn or newborn baby if you take TRINTELLIX during pregnancy.

- Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with TRINTELLIX.

- There is a pregnancy registry for females who are exposed to TRINTELLIX during pregnancy. The purpose of the registry is to collect information about the health of females exposed to TRINTELLIX and their baby. If you become pregnant during treatment with TRINTELLIX, talk to your healthcare provider about registering with the National Pregnancy Registry for Antidepressants at 1-844-405-6185 or visit online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/antidepressants/>.

- are breastfeeding or plan to breastfeed. It is not known if TRINTELLIX passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with TRINTELLIX.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

TRINTELLIX and some other medicines may affect each other causing possible serious side effects. TRINTELLIX may affect the way other medicines work and other medicines may affect the way TRINTELLIX works.

Especially tell your healthcare provider if you take:

- medicines used to treat migraine headache called triptans
- tricyclic antidepressants
- lithium
- tramadol, fentanyl, meperidine, methadone, or other opioids
- tryptophan
- buspirone
- St. John's Wort
- medicines that can affect blood clotting such as aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), warfarin
- diuretics
- medicines used to treat mood, anxiety, psychotic or thought disorders, including selective serotonin reuptake inhibitors (SSRIs) and serotonin norepinephrine reuptake inhibitors (SNRIs)
- medicines used to treat seizures or convulsions

Ask your healthcare provider if you are not sure if you are taking any of these medicines. Your healthcare provider can tell you if it is safe to take TRINTELLIX with your other medicines.

Do not start or stop any other medicines during treatment with TRINTELLIX without talking to your healthcare provider first. Stopping TRINTELLIX suddenly may cause you to have serious side effects. See, "**What are the possible side effects of TRINTELLIX?**"

Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

How should I take TRINTELLIX?

- Take TRINTELLIX exactly as your healthcare provider tells you to take it. Your healthcare provider may need to change the dose of TRINTELLIX until it is the right dose for you.
- Take TRINTELLIX 1 time each day.
- Take TRINTELLIX with or without food.
- If you take too much TRINTELLIX, call your Poison Help Line at 1-800-222-1222 or go to the nearest hospital emergency room right away.

What are the possible side effects of TRINTELLIX?

TRINTELLIX may cause serious side effects, including:

- See, “What is the most important information I should know about TRINTELLIX?”
- **Serotonin syndrome.** A potentially life-threatening problem called serotonin syndrome can happen when you take TRINTELLIX with certain other medicines. See, “Do not take TRINTELLIX if you”. **Call your healthcare provider or go to the nearest hospital emergency room right away** if you have any of the following signs and symptoms of serotonin syndrome:
 - agitation
 - seeing or hearing things that are not real (hallucinations)
 - confusion
 - coma
 - fast heart-beat
 - changes in blood pressure
 - dizziness
 - sweating
 - flushing
 - high body temperature (hyperthermia)
 - shaking (tremors), stiff muscles, or muscle twitching
 - loss of coordination
 - seizures
 - nausea, vomiting, diarrhea
- **Increased risk of bleeding.** Taking TRINTELLIX with aspirin, non-steroidal anti-inflammatory drugs (NSAIDs), warfarin or blood thinners may add to this risk. Tell your healthcare provider right away about any unusual bleeding or bruising.
- **Manic episodes.** Manic episodes may happen in people with bipolar disorder who take TRINTELLIX. Symptoms may include:
 - greatly increased energy
 - racing thoughts
 - unusually grand ideas
 - talking more or faster than usual
 - severe problems sleeping
 - reckless behavior
 - excessive happiness or irritability
- **Discontinuation syndrome.** Suddenly stopping TRINTELLIX may cause you to have serious side effects. Your healthcare provider may want to decrease your dose slowly. Symptoms may include:

◦ nausea	◦ anxiety
◦ sweating	◦ confusion
◦ changes in your mood	◦ headache
◦ irritability and agitation	◦ tiredness
◦ dizziness	◦ problems sleeping
◦ electric shock feeling (paresthesia)	◦ hypomania
◦ tremor	◦ ringing in your ears (tinnitus)
	◦ seizures
- **Eye problems (angle-closure glaucoma).** TRINTELLIX may cause a type of eye problem called angle-closure glaucoma in people with certain other eye conditions. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are. Call your healthcare provider if you have eye pain, changes in your vision, or swelling or redness in or around the eye.
- **Low levels of sodium in your blood (hyponatremia).** Low sodium levels in your blood that may be serious and may cause death, can happen during treatment with TRINTELLIX. Elderly people and people who take certain medicines may be at a greater risk for developing low sodium levels in your blood. Signs and symptoms may include:
 - headache
 - difficulty concentrating
 - memory changes
 - confusion
 - weakness and unsteadiness on your feet, which can lead to falls

In more severe or more sudden cases, signs and symptoms include:

- seeing or hearing things that are not real (hallucinations)
- fainting
- seizures
- coma
- stopping breathing (respiratory arrest)
- **Sexual problems (dysfunction).** Taking TRINTELLIX may cause sexual problems.
Symptoms in males may include:
 - delayed ejaculation or inability to have an ejaculation
 - problems getting or keeping an erection
 - decreased sex driveSymptoms in females may include:
 - decreased sex drive
 - delayed orgasm or inability to have an orgasmTalk to your healthcare provider if you develop any changes in your sexual function or if you have any questions or concerns about sexual problems during treatment with TRINTELLIX. There may be treatments your healthcare provider can suggest.

The most common side effects of TRINTELLIX include nausea, constipation, and vomiting.

These are not all the possible side effects of TRINTELLIX.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store TRINTELLIX?

- Store TRINTELLIX at room temperature between 68°F to 77°F (20°C to 25°C).
- **Keep TRINTELLIX and all medicines out of the reach of children.**

General information about the safe and effective use of TRINTELLIX.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not take TRINTELLIX for a condition for which it was not prescribed. Do not give TRINTELLIX to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about TRINTELLIX that is written for health professionals.

What are the ingredients in TRINTELLIX?

Active ingredient: vortioxetine hydrobromide

Inactive ingredients: mannitol, microcrystalline cellulose, hydroxypropyl cellulose, sodium starch glycolate, magnesium stearate and film coating consisting of hypromellose, titanium dioxide, polyethylene glycol 400, iron oxide red (5 mg and 20 mg) and iron oxide yellow (10 mg)

Distributed and Marketed by:

Takeda Pharmaceuticals America, Inc. Lexington, MA 02421

Marketed by:

Lundbeck, Deerfield, IL 60015

TRINTELLIX is a trademark of H. Lundbeck A/S registered with the U.S. Patent and Trademark Office and used under license by Takeda Pharmaceuticals America, Inc.

©2013-2023 Takeda Pharmaceuticals America, Inc.

For more information, go to www.TRINTELLIX.com or call 1-877-TAKEDA-7 (1-877-825-3327).

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: 8/2023

SPI-0456